

# Discrete Mathematics --- Day 31 --- November 8, 2004

Ernest Byrd


**Binomial Theorem:**

For a natural number n:

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad \text{--Sigma notation means plug in various values of } k \text{ from } 0 \text{ to } n \text{ and add up the terms.}$$

Eg:  $(a+b)^3 = (a+b)(a+b)(a+b)$   
 $= (a+b)(a^2+2ab+b^2)$   
 $= (a^3+2a^2b+ab^2+a^2b+2ab^2+b^3)$   
 $= a^3+3a^2b+3ab^2+b^3$

A good way to show this is by using Pascal's Triangle:



Provided by The Math Forum - <http://forum.swarthmore.edu>

Eg:

$$1$$

$$a + b$$

$$a^2 + 2ab + b^2$$

$$a^3 + 3a^2b + 3ab^2 + b^3$$

.....

This shows how Pascal's Triangle is relevant here.

**Combinational Proof:**

Consider  $(a+b)^n = (a+b)(a+b)(a+b)(a+b)(a+b)\dots(a+b)(a+b)$   
 n Factors

**Here we show it in an “anagram” type of way:**

Consider multiplying 3 binomials together which are not all the same:

$$(a+b)(c+d)(e+f)$$

$$=(ac+ad+bc+bd)(e+f)$$

$$=ace+ade+be+bde+acf+adf+bcf+bdf$$

We see that the result is the sum of all ways to make a product by taking one term from each factor

So, consider:

$$(a+b)^6 = (a+b)(a+b)(a+b)(a+b)(a+b)(a+b)$$

Let’s multiply this out “all at once.” The product will consist of terms of the form  $const \times a^i b^{6-i}$ , where the constant is obtained by grouping together all the terms which have  $i$  "a" factors and  $6-i$  "b" factors. That constant is just the number of anagrams of a word with  $i$  "a"s and  $6-i$  "b"s.

To find out what row we can find our answer on in Pascal' s Triangle we find out how many “anagrams” this would have.

Eg:

$$\frac{6!}{2!4!} = 15$$

Therefore, row 6 of Pascal' s Triangle would give us our answer which is:

$$a^6 + 6a^5b + 15a^4b^2 + 20a^3b^3 + 15a^2b^4 + 6ab^5 + b^6$$