

1 Propositional Logic

1.1 Syntax of propositional logic

Rosen and Zybooks use term *compound proposition* for an expression written using logic. I will use the more common term *propositional formula* instead.

The *syntax* of propositional logic only says what a propositional formula looks like. It does not say what a propositional formula means. We use A , B , C and ϕ (Greek phi) to name arbitrary propositional formulas.

Definition 1.1. A *propositional formula* is defined as follows.

1. Symbols \mathbf{T} and \mathbf{F} are propositional formulas.
2. A *propositional variable* is a propositional formula. We will use p , q , r and s , possibly with subscripts, as propositional variables and X for talking about an arbitrary variable.
3. If A and B are propositional formulas then so are
 - (a) $A \vee B$,
 - (b) $A \wedge B$,
 - (c) $\neg A$,
 - (d) (A) .

For example, each of the following is a propositional formula.

- p
- $p \vee q$
- $p \wedge \neg q$
- $p \wedge q \wedge r$

- $q \vee p \wedge r$
- $(r \wedge \mathbf{T}) \vee \neg q$

Operator \vee is read “or”, \wedge is read “and”, and \neg is read “not”.

Rules of *precedence* and *associativity* determine how you break a propositional formula into subformulas. Higher precedence operators are done first. The following lists operators by precedence, from highest to lowest.

Precedence	
parentheses	high
\neg	
\wedge	
\vee	low

For example, $p \vee q \wedge r$ is understood to have the same structure as $p \vee (q \wedge r)$ since \wedge has higher precedence than \vee .

Associativity determines how an expression is broken into subexpressions when it involves two or more occurrences of the same operator. We assume that operators \vee and \wedge are done from left to right. That is, they are *left-associative*. (Associativity is like the wind. A north wind blows from north to south.) For example, $p \vee q \vee r$ has the same structure as $(p \vee q) \vee r$. Associativity does not really matter for \vee and \wedge because they are *associative operators*. That is, $(p \vee q) \vee r$ always has the same meaning as $p \vee (q \vee r)$ and $(p \wedge q) \wedge r$ always has the same meaning as $p \wedge (q \wedge r)$. But associativity does matter for some operators, so it is wise to think about it.

1.2 Meaning of propositional logic

The meaning of a propositional formula can only be defined when the values of all of its variables are given. Each variable can be true or false.

Definition 1.2. A *truth-value assignment* is a set of components of the form $X = V$ where X is a variable and V is either T or F. For example,

$\{p=T, q=F\}$ is a truth-value assignment. (Note that T and F are possible values of a propositional variable or a propositional formula. Do not confuse them with **T** and **F**, which are propositional formulas.)

Definition 1.3. If a is a truth-value assignment and X is a variable then $a(X)$ is the value (T or F) that a gives for variable X . For example, if a is $\{p=T, q=F\}$ then $a(p) = T$ and $a(q) = F$.

Definition 1.4. Suppose that ϕ is a propositional formula and a is a truth-value assignment that defines every variable that occurs in ϕ . Notation $\phi : a$ indicates the value of ϕ (either T or F) when variables have values given by a . Specifically:

1. $(\mathbf{T} : a) = T$. That is, symbol **T** is always true; it does not depend on a .
2. $(\mathbf{F} : a) = F$. That is, symbol **F** is always false; it does not depend on a .
3. If X is a variable then $(X : a) = a(X)$. That is, X has the value that it is given by truth-value assignment a .
4. $(A \vee B : a)$ is T if at least one of $(A : a)$ and $(B : a)$ is T, and is F otherwise. For example, $((p \vee q) : \{p = T, q = F\})$ is T because $(p : \{p = T, q = F\})$ is T, and we only need one of p and q to be true.
5. $((A \wedge B) : a)$ is T if both of $(A : a)$ and $(B : a)$ are T, and is F otherwise. For example, $((p \wedge q) : \{p = T, q = F\})$ is F because $(p : \{p = T, q = F\})$ and $(q : \{p = T, q = F\})$ are not both T.
6. $(\neg A : a)$ is T if $(A : a)$ is F, and is F if $(A : a)$ is T.
7. $((A) : a) = (A : a)$. Parentheses only influence the structure of a propositional formula. A parenthesized formula (A) has the same meaning as A .

You determine the value of a propositional formula by building up larger and larger subexpressions, being careful to follow the rules of precedence and associativity. For example, suppose that $a = \{p=F, q=T, r=T\}$. Then

- (a) $(q : a) = \text{T}$
- (b) $(p : a) = \text{F}$
- (c) $(\neg p : a) = \text{T}$ by (b)
- (d) $(\neg p \wedge q : a) = \text{T}$ by (a) and (c)

1.3 Additional definitions

Definition 1.5. $A \rightarrow B$ is defined to be an abbreviation for $\neg A \vee B$. Operator \rightarrow is read “implies”.

Intuitively, $A \rightarrow B$ means “if A is true then B is true.” But that is not its definition. Its definition is that either A is false or B is true (or both). Notice that, if B is true, then $A \rightarrow B$ is true, *by definition*. Also, if A is false, then $A \rightarrow B$ is true, *by definition*.

Operator \rightarrow has lower precedence than \vee and is left-associative. Note that \rightarrow is not an associative operator. $(A \rightarrow B) \rightarrow C$ does not have the same meaning as $A \rightarrow (B \rightarrow C)$.

Definition 1.6. $A \leftrightarrow B$ is defined to be the same as $((A \rightarrow B) \wedge (B \rightarrow A))$. Operator \leftrightarrow is read “if and only if”.

Formula $A \leftrightarrow B$ says that A and B have the same value; either both are true or both are false. In fact, $A \leftrightarrow B$ is equivalent to $(A \wedge B) \vee (\neg A \wedge \neg B)$. That is, either A and B are both true or A and B are both false.

Definition 1.6. $A \equiv B$ if $A \leftrightarrow B$ is true for all possible values of the variable in A and B . Read $A \equiv B$ as “ A is equivalent to B .”

Operators \leftrightarrow and \equiv have even lower precedence than \rightarrow . Here is a complete precedence table, from high to low precedence.

Precedence	
parentheses	high
\neg	
\wedge	
\vee	
\rightarrow	
\leftrightarrow	
\equiv	low

1.4 Truth tables

Since the value of a propositional formula depends on the values of its variables, one way to understand what the formula means is to look at its value for all possible values of the variables. That leads to the idea of a *truth table* of a propositional formula. The following is a truth table for $\neg p \vee q$.

p	q	\neg	p	\vee	q
F	F	T	F	T	F
F	T	T	F	T	T
T	F	F	T	F	F
T	T	F	T	T	T

There is a row for each possible collection of values of the variables. Under each variable, we write that variable's value. Under each operator, we write the value of the formula having that operator as its main or outermost operator. The column in blue is the value of the entire formula, $\neg p \vee q$.

1.5 Validity

Definition 1.8. Propositional formula ϕ is *valid* if $(\phi : a)$ is true for every truth value assignment a . A valid formula is also called a *tautology*.

For example, operator \vee is commutative. Another way to say that is to say that formula

$$(p \vee q) \leftrightarrow (q \vee p)$$

is valid. Let's check that using a truth table.

p	q	$(p \vee q)$	\leftrightarrow	$(q \vee p)$
F	F	F	T	F
F	T	T	T	T
T	F	T	T	T
T	T	T	T	T

The validity of

$$(p \vee q) \leftrightarrow (q \vee p)$$

is evident from the blue column of all Ts.

Table 1-1 shows a collection of true equivalences and valid propositional formulas. You can check each one using a truth table.

Valid equivalences give you a way to replace one formula by another. For example, if you see $p \vee q$ in any context, you can replace it by $q \vee p$. In fact, you can replace any variable by any propositional formula in any of the above tautologies (or any other valid propositional formula) and they are still valid, provided (1) you replace every occurrence of a variable by the same propositional formula and (2) you use parentheses to avoid rules of precedence from rearranging the formula. For example, the commutative law for \wedge says that

$$p \wedge q \equiv q \wedge p.$$

Replacing p by $(w \rightarrow v)$ and q by $\neg r$ yields

$$(w \rightarrow v) \wedge \neg r \equiv \neg r \wedge (w \rightarrow v)$$

which is also valid.

Table 1-1: Some propositional tautologies	
Equivalence	Name
$\neg(\neg p) \equiv p$	double negation
$p \vee q \equiv (q \vee p)$	commutative law of \vee
$p \wedge q \equiv (q \wedge p)$	commutative law of \wedge
$(p \vee q) \vee r \equiv p \vee (q \vee r)$	associative law of \vee
$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$	associative law of \wedge
$(p \wedge (q \vee r)) \equiv (p \wedge q) \vee (p \wedge r)$	distributive law of \wedge over \vee
$(p \vee (q \wedge r)) \equiv (p \vee q) \wedge (p \vee r)$	distributive law of \vee over \wedge
$\neg(p \vee q) \equiv \neg p \wedge \neg q$	DeMorgan's law for \vee
$\neg(p \wedge q) \equiv \neg p \vee \neg q$	DeMorgan's law for \wedge
$\neg(p \rightarrow q) \equiv p \wedge \neg q$	DeMorgan's law for \rightarrow
$p \rightarrow q \equiv \neg q \rightarrow \neg p$	Law of the contrapositive
$(p \vee q) \rightarrow r \equiv (p \rightarrow r) \wedge (q \rightarrow r)$	cases
$(p \wedge q) \rightarrow r \equiv (p \rightarrow (q \rightarrow r))$	
$p \wedge \neg p \equiv \mathbf{F}$	contradiction 1
$p \equiv (\neg p \rightarrow p)$	contradiction 2
$p \equiv (\neg p \rightarrow \mathbf{F})$	contradiction 3
$p \vee \neg p$	Law of the excluded middle
$p \rightarrow p$	Law of the excluded middle, re-stated using \rightarrow
$\neg(p \wedge \neg p)$	Law of the excluded middle (De-Morgan variant)
$p \rightarrow (q \rightarrow p)$	
$\neg p \rightarrow (p \rightarrow q)$	

next